

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁴ : C07H 21/04, C12N 1/20, 15/00 C07K 13/00	A1	(11) International Publication Number: WO 89/ 09223 (43) International Publication Date: 5 October 1989 (05.10.89)
(21) International Application Number: PCT/US89/01238 (22) International Filing Date: 24 March 1989 (24.03.89) (31) Priority Application Numbers: 176,107 253,807 312,763 (32) Priority Dates: 30 March 1988 (30.03.88) 5 October 1988 (05.10.88) 21 February 1989 (21.02.89) (33) Priority Country: US (71) Applicant: ARCH DEVELOPMENT CORPORATION [US/US]; 1115-25 East 58th Street, Chicago, IL 60637 (US). (72) Inventors: LIAO, Shutsung ; 5632 South Woodlawn Avenue, Chicago, IL 60637 (US). CHANG, Chawn-shang ; 5644 South Harper Avenue, Chicago, IL 60637 (US).		(74) Agent: MEINERT, M., C.; Marshall, O'Toole, Gers- tein, Murray & Bicknell, Two First National Plaza, Chicago, IL 60603 (US). (81) Designated States: AT (European patent), BE (Euro- pean patent), CH (European patent), DE (European patent), FR (European patent), GB (European pa- tent), IT (European patent), JP, LU (European pa- tent), NL (European patent), SE (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: DNA BINDING PROTEINS INCLUDING ANDROGEN RECEPTOR (57) Abstract Disclosed are DNA sequences encoding DNA binding polypeptides including androgen receptor (AR) and TR2 polypeptides. Illustratively, human and rat AR-cDNA have 79 kD and 98 kD polypeptide expression products which are immunoprecipitable by human auto-immune anti-androgen receptor antibodies and are capable of binding androgens specifically and with high affinity. Also disclosed are antibodies and immunological methods and materials for detection of androgen receptor and TR2 polypeptides and hybridization methods and materials for detection of AR and TR2-related nucleic acids.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

- 1 -

"DNA BINDING PROTEINS INCLUDING ANDROGEN RECEPTOR"

5

10

BACKGROUND OF THE INVENTION

15 The present invention relates generally to DNA binding regulatory proteins and more particularly to DNA sequences encoding androgen receptor protein and novel DNA binding proteins designated TR2, to the polypeptide products of recombinant expression of these DNA sequences, to peptides whose sequences are based on amino acid sequences deduced from these DNA sequences, 20 to antibodies specific for such proteins and peptides, and to procedures for detection and quantification of such proteins and nucleic acids related thereto.

25 There are five major classes of steroid hormones: progestins, glucocorticoids, mineralocorticoids, androgens, and estrogens. Receptor proteins, each specific for a steroid hormone, are distributed in a tissue specific fashion and in target cells, steroid hormones can form specific complexes with corresponding intracellular receptors. [Jensen, et al., Proc. Nat'l. Acad. Sci. (USA), 59:632 (1968); Gorski, et al., Ann. Rev. Physiol., 38:425-450 (1976); and Liao, et al., page 633 in Biochemistry of Hormones, H.L.J. Makin, ed. (Blackwell Sci. Publ. Oxford, 1984)]. The hormonal 30 regulation of gene expression appears to involve interaction of steroid receptor complexes with certain segments of genomes and modulation of specific gene trans- 35

- 2 -

cription. See, e.g., Ringold, Ann. Rev. Pharmacol. Toxicol., 25:529 (1985); and Yamamoto, Ann. Rev. Genet., 19:209 (1985). Many of the primary effects of hormones involve increased transcription of a subset of genes in specific cell types.

The successful cloning of e.g., cDNAs coding for various steroid receptors has allowed the structural and functional analysis of different steroid receptor domains involved in steroid and DNA binding. See, e.g., Hollenberg, et al., Nature (London), 318:635 (1985); Miesfeld, et al., Cell, 46:389 (1986); Danielsen, et al., EMBO J., 5:2513 (1986); Greene, et al., Science, 231:1150 (1986); Green, et al., Nature (London), 320:134 (1986); Krust, et al., EMBO J., 5:891 (1986); Loosfelt, et al., Proc. Nat'l. Acad. Sci. (USA), 83:9045 (1986); Conneely, et al., Science, 233:767 (1987); Law, et al., Proc. Nat'l. Acad. Sci. (USA), 84:2877 (1987); Misrahi, et al., Biochem. Biophys. Res. Commun., 143:740 (1987); Arriza, et al., Science, 237:268 (1987); Sap, et al., Nature (London), 324:635 (1986); Weinberger, et al., Nature (London), 318:641 (1986); Benbrook, et al., Science, 238:788 (1987); and Evans, Science, 240:889 (1988).

Androgens, such as testosterone, are responsible for the development of male secondary sex characteristics and are synthesized primarily in testis. Cloning of a cDNA for androgen receptor (AR) has been difficult because, until recently, monospecific antibodies against AR have not been available for screening cDNA libraries. An abstract by Govindan, et al., J. Endocrinol. Invest., 10 (Suppl. 2) (1987), reported the isolation of cDNA clones encoding human androgen receptor isolated from a human testis λ gt-11 cDNA library using synthetic oligonucleotides homologous to human glucocorticoid, estradiol, and progesterone receptors as probes. The expressed protein reportedly

- 3 -

bound tritium-labelled DHT (dihydrotestosterone) with high affinity and specificity. However, no nucleotide or amino acid sequence analysis was provided for full length androgen receptors, nor was any description provided concerning isolation of the full length putative androgen receptor clones.

Recently, Chang, C., et al., Science, 240:324 (April 15, 1988), co-authored by the inventors herein, described cDNAs encoding androgen receptors obtained from human testis and rat ventral prostate cDNA libraries. These cDNAs for human and rat androgen receptor were reported to be long enough to code for 94 kDa and 76 kDa receptors. The molecular weights were derived with the assistance of a software program known as: DNA Inspector II (Textco West Lebanon, New Hampshire) open reading frame analysis. With a new DNA Inspector IIe program, hAR (918 amino acids) has an estimated molecular weight 98,608 and rAR (902 amino acids) has a molecular weight of 98,133. Therefore, the reported "94 kDa" AR is now termed "98 kDa" AR; and the hAR or rAR polypeptides, from the second ATG/Met, reported as "76 kDa" are now termed "79 kDa". See also, Chang, C., et al., Proc. Nat'l. Acad. Sci. (USA), 85:7211 (October 5, 1988) also co-authored by the inventors herein.

In contrast, Lubahn, D., et al., Science, 240:327 (1988), using libraries from human epididymis and cultured human foreskin fibroblasts obtained a human cDNA which was expressed in monkey kidney (COS) cells to yield a protein, present in the cytosol, capable of binding androgens. This cDNA, however, was only sufficient to code for a receptor having an estimated molecular weight of 41,000. Therefore, the cDNA obtained only coded for a portion of AR.

- 4 -

Of interest to the present invention is Young, et al., Endocrinol., 123:601 (1988), wherein the production of anti-AR monoclonal antibodies was reported. Anti-AR autoantibodies were identified in the sera of prostate cancer patients, as described in Liao, S., et al., Proc. Nat'l. Acad. Sci. (USA), 82:8345 (1984) (one of the co-inventors herein), and were characterized with respect to their titer, affinity, and specificity. Subsequently, lymphocytes from the blood of those patients having high antibody titers were isolated, transformed with Epstein-Barr Virus (EBV), and cloned for anti-AR monoclonal antibody production. These monoclonal antibodies were found to interact with androgen receptors from rat prostate. An attempt to scale-up antibody production resulted in a decline of antibody secretion. It is not uncommon for transformed B-cells to be more unstable than hybridoma cells. Kozbor, et al., Eur. J. Immunol., 14, 23 (1984). Because of the instability associated with such cell lines, an alternate source of monoclonal antibodies is preferred.

There thus exists a need in the art for information concerning the primary structural conformation of androgen receptor protein and other DNA binding proteins such as might be provided by knowledge of human and other mammalian DNA sequences encoding the same. Availability of such DNA sequences would make possible the application of recombinant methods to the large scale production of the proteins in procaryotic and eukaryotic host cells, as well as DNA-DNA, DNA-RNA, and RNA-RNA, hybridization procedures for the detection, quantification and/or isolation of nucleic acids associated with the proteins. Possession of androgen receptor and related DNA-binding proteins and/or knowledge of the amino acid sequences of the same would make possible, in turn, the development of monoclonal and

- 5 -

polyclonal antibodies thereto (including antibodies to protein fragments or synthetic peptides modeled thereon) for the use in immunological methods for the detection and quantification of the proteins in fluid and tissue samples as well as for tissue specific delivery of substances such as labels and therapeutic agents to cells expressing the proteins.

BRIEF SUMMARY OF THE INVENTION

10

The present invention provides novel purified and isolated DNA sequences encoding androgen receptor protein and a structurally related protein, designated TR2 protein, which also has DNA binding (and hence DNA replication or transcription regulatory) capacity. In presently preferred forms, novel DNA sequences comprise cDNA sequences encoding human and rat androgen receptor and human TR2 protein. Alternate DNA forms such as genomic DNA, and DNA prepared by partial or total chemical synthesis from nucleotides as well as DNA with deletions or mutations, is also within the contemplation of the invention.

Association of DNA sequences provided by the invention with homologous or heterologous species expression control DNA sequences, such as promoters, operators, regulators and the like, allows for in vivo and in vitro transcription to form messenger RNA which, in turn, is susceptible to translation to provide androgen receptor and TR2 proteins, and related poly- and oligo-peptides in large quantities. In a presently preferred DNA expression system of the invention, AR and TR2 encoding DNA is operatively associated with a viral (T7) regulatory (promoter) DNA sequence allowing for in vitro transcription and translation in a cell free system to provide, e.g., a 79 kD and 98 kD human androgen receptor (hAR) protein, 79 kD and 98 kD rat

- 6 -

androgen receptor (rAR) protein and smaller forms of these proteins, as well as TR2 protein, including 20 kD and 52 kD species.

5 Incorporation of DNA sequences into procaryotic and eucaryotic host cells by standard transformation and transfection processes, potentially involving suitable viral and circular DNA plasmid vectors, is also within the contemplation of the invention and is expected to provide useful proteins in quantities heretofore unavailable from natural sources. Systems provided by the invention included transformed E. coli DH5 α cells, deposited January 25, 1989, with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852 in accordance with the U.S. Patent and Trademark Office's requirements for microorganism deposits, and designated EC-hAR3600 under A.T.C.C. Accession No. 67879; EC-rAR 2830, A.T.C.C. No. 67878; EC-TR2-5, A.T.C.C. No. 67877; and EC TR2-7, A.T.C.C. No. 67876. Use of mammalian host cells is expected to provide for such post-translational modifications (e.g., truncation, glycosylation, and tyrosine, serine or threonine phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention.

25 Novel protein products of the invention include polypeptides having the primary structural conformation (i.e., amino acid sequence) of AR and TR2 proteins as well as peptide fragments thereof and synthetic peptides assembled to be duplicative of amino acid sequences thereof. Proteins, protein fragments, and synthetic peptides of the invention are projected to have numerous uses including therapeutic, diagnostic and prognostic uses and will provide the basis for preparation of monoclonal and polyclonal antibodies specifically immunoreactive with AR and TR2 proteins. Preferred protein fragments and synthetic peptides

- 7 -

include those duplicating regions of AR and TR2 proteins which are not involved in DNA binding functions and the most preferred are those which share at least one antigenic epitope with AR and TR2 proteins.

5 Also provided by the present invention are polyclonal and monoclonal antibodies characterized by their ability to bind with high immunospecificity to AR and TR2 proteins and to their fragments and peptides, recognizing unique epitopes which are not common to
10 other proteins especially DNA binding proteins.

Illustratively provided according to the present invention are monoclonal antibodies, designated AN1-6, AN1-7, AN1-15; and produced by hybridoma cell lines designated H-AN1-6, H-AN1-7, H-AN1-15; deposited
15 January 25, 1989, under Accession Nos. HB 10,000; HB 9,999; and HB 10,001, respectively, with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852 in accordance with the U.S. Patent and Trademark Office's requirements for
20 microorganism deposits. These antibodies are characterized by (a) capacity to bind androgen receptors from rat ventral prostate and synthetic peptides having sequences predicted from the structure of hAR-cDNA and rAR-cDNA; (b) specific immunological reactivity with,
25 and capacity to reversibly immunobind to, naturally occurring and recombinant androgen receptors, in native and denatured conformations; and (c) specific immunological reactivity with, and capacity to reversibly immunobind to, proteinaceous materials
30 including all or a substantially, immunologically significant, part of an amino acid sequence duplicative of that extant at residues 331 through 577 of hAR and corresponding amino acid sequences in rAR.

The monoclonal antibodies of the invention can
35 be used for affinity purification of AR from human or rat prostate, and other sources such as AR-rich organs and cultured cells.

- 8 -

Also provided by the present invention are novel procedures for the detection and/or quantification of normal, abnormal, or mutated forms of AR and TR2, as well as nucleic acids (e.g., DNA and mRNA) associated therewith. Illustratively, antibodies of the invention may be employed in known immunological procedures for quantitative detection of AR and TR2 proteins in fluid and tissue samples, of DNA sequences of the invention (particularly those having sequences encoding DNA binding proteins) that may be suitably labelled and employed for quantitative detection of mRNA encoding these proteins.

Among the multiple aspects of the present invention, therefore, is the provision of (a) novel AR and TR2-encoding DNA sequences set out in Figure 3, as well as (b) AR and TR2-encoding DNA sequences which hybridize thereto under hybridization conditions of the stringency equal to or greater than the conditions described herein and employed in the initial isolation of cDNAs of the invention, and (c) DNA sequences encoding the same allelic variant, or analog AR and TR2 polypeptides through use of, at least in part, degenerate codons. Correspondingly provided are viral or circular plasmid DNA vectors incorporating such DNA sequences and procaryotic and eucaryotic host cells transformed or transfected with such DNA sequences and vectors as well as novel methods for the recombinant production of AR and TR2 proteins through cultured growth of such hosts and isolation of these proteins from the hosts or their culture media.

Preferred polypeptide products of the invention include the approximately 79 kD (starting from the second ATG/Met) and 98 kD (starting from the first ATG/Met) hAR polypeptides having the deduced amino acid sequence of 734 and 918 residues, respectively, as set out in Figure 3. Also preferred are the 79 kD and 98 kD

- 9 -

rAR species polypeptides having the deduced sequence of 733 and 902 residues set out in Figure 3 and the 20 kD and 52 kD species human TR2 polypeptides having the same deduced amino acid sequence of 184 and 483 residues set out in Figure 4. The preferred 79 kD and 98 kD hAR and rAR polypeptides may be produced in vitro and are characterized by a capacity to specifically bind androgens with high specificity and by their immunoprecipitatability by human auto-immune anti-androgen receptor antibodies.

Other aspects and advantages of the present invention will be apparent upon consideration of the following detailed description thereof which includes numerous illustrative examples of the practice of the invention, reference being made to the drawing wherein:

Figure 1 illustrates the strategy employed in construction of a human androgen receptor cDNA vector;

Figure 2 illustrates the strategy employed in construction of rat androgen receptor cDNA vectors;

Figure 3 provides a 3715 base pair nucleotide sequence for a human androgen receptor (hAR) DNA clone and the deduced sequence of 734 and 918 amino acid residues for hAR proteins; and in addition provides a 3218 base pair nucleotide sequence for a rat androgen receptor (rAR) DNA clone and the deduced sequences of 733 and 902 amino acids for two rAR species;

Figure 4 provides a 2029 base pair nucleotide sequence for a human TR2 DNA clone and a deduced sequence of 483 amino acids for a "TR2-5" species and a deduced sequence of 184 amino acids for a "TR2-7" species; and

Figure 5 provides an amino acid sequence alignment of the cysteine-rich DNA binding domain of human androgen receptor, glucocorticoid receptor, mineralocorticoid receptor, progesterone receptor, estrogen receptor, TR2, rat AR, chick vitamin D receptor

- 10 -

(c-VDR), and the v-erb A oncogene product of avian erythroblastosis virus.

Figures 6, 7, and 8 illustrate, respectively, the in-frame fusion of three different parts of the AR gene (the N-terminal, the DNA-binding domain and the androgen-binding domain) to the N-terminal half of the trpE gene using pATH expression vectors.

DETAILED DESCRIPTION

10

The following examples illustrate practice of the invention. Example 1 relates to the isolation, preparation, and partial structural analysis of cDNA for human and rat androgen receptors. Example 2 relates to confirmation of the presence on the human X-chromosome of an AR-type cDNA sequence. Example 3 relates to the preparation of human and rat cDNAs containing AR-type cDNA from different clones and ligation into the pGEM-3Z plasmid. Example 4 relates to transcription and translation of the AR-type cDNA plasmid DNA. Example 5 relates to steroid binding activity of the expression product of Example 4. Example 6 relates to the binding activity of the expression product of Example 4 to human auto-antibodies. Example 7 relates to the characterization of TR2-cDNA. Example 8 relates to the in vitro transcription and translation of TR2-cDNA. Example 9 relates to the binding activity of TR2-cDNA expression product. Example 10 relate to the androgen regulation of TR2 mRNA levels in the rat ventral prostate. Example 11 relates to recombinant expression systems of the invention. Example 12 relates to the production of fusion proteins and their use in producing polyclonal and monoclonal antibodies according to the invention. Example 13 relates to use of DNA probes of the inventions. Example 14 relates to development of transgenic animals by means of DNA sequences of the invention.

- 11 -

These examples are for illustrative purposes only and are not intended in any way to limit the scope of the invention.

5

EXAMPLE 1Preparation and Partial Structural Analysis of cDNA for Human and Rat Androgen Receptors

10 The isolation of cDNA for human androgen receptor (hAR) and rat androgen receptor (rAR) was accomplished using λ GT11 cDNA libraries. The human testis and prostate λ GT11 libraries were obtained from Clontech Co., Palo Alto, California and a rat ventral prostate λ GT11 library in *E.coli* Y1090 was constructed
15 as described in Chang, et al., *J. Biol. Chem.*, 262:11901 (1987). In general, clones were differentiated using oligonucleotide probes specific for various steroid receptors. The cDNA libraries were initially screened with a set of 41-bp oligonucleotide probes designed for
20 homology to nucleotide sequences in the DNA-binding domain of glucocorticoid receptors (GR), estrogen receptors (ER), progesterone receptors (PR), mineralocorticoid receptors (MR), and the v-erb A oncogene product of avian erythroblastosis virus. The set of probes had the
25 following sequence: TGTGGAAGCTGT/CAAAGTC/ATTCTTTAAAAGG/AGCAA/GTGAAGG.

The plaques were replicated on a nitrocellulose filter and screened with a 5'-end 32 P-labeled 41-bp oligonucleotide probes. The conditions of hybridization
30 were 25% formamide, 5X Denhardt's solution (0.1% Ficoll 400, 0.1% polyvinylpyrrolidone, 0.1% bovine serum albumin), 0.1% SDS, 5X SSC (1X SSC is 150 mM NaCl, 15 mM sodium citrate), 100 μ g/ml denatured salmon sperm DNA, and 1 μ g/ml poly(A) at 30°C. Filters were washed with a
35 solution containing 0.1% SDS, 0.05% sodium pyrophosphate and 0.4X SSC at 37°C.

- 12 -

A less stringent hybridization condition (2X SSC at 37°C) was used for the first screen employing the 41 bp probes. The remaining clones were then probed again at more stringent conditions by reducing the concentration of SSC, eventually to 0.4X SSC at 37°C, or by increasing the temperature, or by increasing the concentration of formamide. In some procedures, 5X SSC, 8% dextran sulfate, and 20% formamide, at 42°C was employed and the result was equivalent to that obtained with 0.6X SSC.

From approximately 3×10^6 human testis recombinants and 6×10^5 rat ventral prostate recombinants, 302 and 21 positive clones, respectively, were obtained.

Based on the assumption that AR might have a cysteine-rich DNA binding domain highly homologous to the DNA-binding regions of other steroid receptors, positive clones from the first screenings were probed with 5'-end ^{32}P -labeled 24-bp oligonucleotides specific for the various steroid receptors for the possible presence of cDNA for AR through a process of elimination. The GR-cDNA clones were eliminated by screening with two GR-specific 24-bp probes that had nucleotide sequences identical to nucleotide segments immediately next to the 5'-end or the 3'-end of the DNA binding-region of hGR-cDNA, i.e., TGTAAGCTCTCCTCCATCCAGCTC and CAGCAGGCCACTACAGGAGTCTCA. 244 and 14 clones, respectively, were eliminated as hGR- and rGR-cDNA clones.

Using similar procedures involving four 24-bp probes for the 5'-end of PR(CCGGATTCAGAAA/GCCAGT/-CCAGAGC) and two 24-bp probes for the 3'-end of ER(GCA/-CGACCAGATGGTCAGTGCCTTG), no ER- or PR-cDNA clones were detected in the human testis library. In the rat prostate library, no ER-cDNA clones were detected but one positive clone was obtained with hPR-specific 24 bp probes.

- 13 -

Following this process of eliminating clones putatively encoding other steroid receptors, the DNA inserts in the remaining clones were analyzed by restriction mapping and subcloned into M13 vectors for di-deoxy sequence analysis. See, Chang, et al., J. Biol. Chem., 262:2826 (1987). Nucleotide sequence analysis allowed four clones to be identified as hMR-cDNA clones.

Through this stepwise process of elimination, 54 human testis clones and 6 rat prostate clones were selected and were then categorized into two groups: 30 human testis clones had sequences overlapping to form a 2.1 kb cDNA; and 24 human testis and 6 rat prostate clones had sequences overlapping to form a cDNA of about 2.7 kb. The two groups of cDNA were designated, respectively, as "TR2-type" and "AR-type" cDNA.

EXAMPLE 2

20 Confirmation of the Presence on the Human X-Chromosome of an AR-type cDNA Sequence Rather than a TR2-type cDNA Sequence

The length between the putative polyadenylation signal (AATAAA) and the 5'-end in the "TR-2 type" cDNA is only 2.0 kb, which is considerably shorter than that for the cDNA of other steroid receptors. Therefore, it was suspected that the "AR-type" cDNA, rather than the "TR2-type" cDNA, encoded androgen receptor. To obtain additional information, a human X-chromosome library prepared according to Kunkel, et al., Nucleic Acids Research, 11:7961 (1983) was probed with the TR2-type cDNA and AR-type cDNA of Example 1. With TR2-type cDNA fragments, no positive clones were detected, while 3 positive clones were obtained with a 1.9 kb fragment of AR-type cDNA from a human testis (clone AR 132), thereby confirming the presence of an AR-type cDNA

- 14 -

sequence on the human X-chromosome. Because the X-chromosome has been implicated as the chromosome which contains an AR gene [Lyon, et al., Nature (London), 227:1217 (1970); Meyer, et al., Proc. Nat'l. Acad. Sci. (USA), 72:1469 (1975); and Amrhein, et al., Proc. Nat'l. Acad. Sci. (USA), 73:891 (1976)], this information suggested that "AR-type" cDNA, but probably not the "TR2-type" cDNA, contained the DNA sequence that could encode for androgen receptor.

Two human clones containing DNA inserts that overlapped to form a 2.7 kb cDNA were designated AR 132 and AR 5. Two rat clones containing DNA inserts that overlapped to form a 2.8 kb cDNA were designated rAR 1 and rAR 4. After restriction enzyme digestion, the DNA segments from these AR-type clones were ligated, selected and amplified using pBR322 and pGEM-3Z vectors as described in Example 3 below.

EXAMPLE 3

20

A. Preparation of a Human cDNA Containing AR-type cDNA from Two Different Clones and Ligation Into the Cloning Vector pGEM-3Z Plasmid

Figure 1 relates to the strategy employed in the construction of a full length hAR-cDNA clone. cDNA of clone AR 132 was digested with Eco RI to obtain a 1.9 kb fragment which was then digested with Kpn I to provide a 1 kb Eco RI-Kpn I fragment. This 1 kb fragment was ligated to a 3 kb fragment obtained by digestion of clone AR 5 with Kpn I and Pvu I. The resulting 4 kb fragment was inserted into Eco RI and Pvu I-digested pBR322 vector and used to infect E. coli DH5 α . The transformed clones were selected by tetracycline-resistance. The plasmid with the DNA insert was digested with Cla I and Nde I to obtain a 2.6 kb fragment. The fragment was blunt-ended with the Klenow fragment of E.

- 15 -

coli DNA polymerase I and ligated to the cloning vector pGEM-3Z plasmid DNA (Promega Biotec, Madison WI.) which was previously blunt-ended by digestion with Sma I. E. coli DH5 α cells were transformed with the plasmid so formed (designated plasmid PhAR3600) and colonies containing the plasmid were selected by ampicillin resistance and amplified. E. coli DH5 α cells, transformed with plasmid PhAR3600, were designated EC-hAR3600 and were deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852 on January 25, 1989 under Accession No. 67879.

The plasmid DNA was isolated and its structure analyzed by restriction enzyme mapping and sequencing. The 2.0 kb hAR fragment obtained by NruI-BamHI digestion of a 2.6 kb hAR in pGEM3Z was then ligated to another 1.6 kb ECORI-NruI fragment of hHR to obtain the full length 3715 bp hAR. The open reading frame is about 2.8 kb which is sufficient to code for a protein with more than 900 amino acids. Near the middle of the protein is a cysteine-rich region with a 72 amino acid sequence highly homologous to regions in other steroid receptors considered to be the DNA binding domain.

As set out in detail below and illustrated in Figure 2, formation of "full length" rat AR clones by slightly varying procedures results in constructions providing RNA transcripts translatable to 79 kD and 98 kD protein products.

B. Preparation of a Rat
2.7 kb cDNA and Ligation
Into the Cloning Vector
pGEM-3Z Plasmid

The 2.4 kb Eco RI-Eco RI cDNA insert of clone rAR 1 was digested with Xmn I to obtain a 2.3 kb fragment. This 2.3 kb Xmn I-EcoR I fragment was ligated to a 400 bp fragment that was obtained by

- 16 -

digestion of another cDNA clone insert (Eco RI-Eco RI insert of rAR 4) with Pst I. The ligated 2.7 kb fragment was inserted into Sma I and Pst I-digested pGEM-3Z vector and used to infect E. coli DH5 α . The E. coli DH5 α cells were transformed with the plasmid and colonies containing the plasmid were selected by ampicillin resistance and amplified. These cells were designated EC-rAR 2830 and were deposited with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852 on January 25, 1989 under Accession No. 67878. As noted in Figure 2, this construction allowed for a transcription product translated beginning with the second of two in-frame methionine-specifying codons (designated ATG₂).

C. Preparation of a Rat
2.83 kb cDNA Ligation
Into the Cloning Vector
pGEM-3Z Plasmid

The 2.4 kb Eco RI-Eco RI cDNA insert of rAR 1 was digested with Hind III to obtain a 1.68 kb fragment. The 1.68 kb Eco RI-Hind III fragment was ligated to a 1.15 kb DNA fragment obtained by digestion of another cDNA clone insert (rAR 6) with Hind III and Pst I. The ligated 2.83 kb fragment was inserted into Eco RI and Pst I-digested pGEM 3Z vector and used to infect E. coli DH5 α . E. coli (DH5 α) cells were transformed with the plasmid and colonies containing the plasmid were selected by ampicillin resistance and amplified. As noted in Figure 2, this construction allowed for a transcription product translated beginning at the first of two in-frame methionine-specifying codons (designated ATG₁).

Figure 3 provides the nucleotide sequence of the DNA sequence of the longer "full length" rat and human AR clones and includes the deduced amino acid sequences. The first and second methionine-specifying

- 17 -

codons are designated at amino acid positions 1 and 170 of rAR and positions 1 and 185 of hAR.

EXAMPLE 4

5

Transcription and Translation of the Human AR-type cDNA Plasmid in a Rabbit Reticulocyte Lysate System

10 pGEM-3Z vector (20 µg) containing 2.6 kb hAR
DNA segment, as described in Example 3, was linearized
with restriction enzyme Bam HI, phenol/chloroform
extracted, and precipitated with ethanol. The
linearized plasmid was transcribed in a reaction mixture
containing 40 mM Tris-HCl, pH 7.5, 6 mM MgCl₂, 2 mM
15 spermidine, 10 mM NaCl, 10 mM DTT, 500 µM each of ATP,
GTP, CTP, and UTP, 160 units ribonuclease inhibitor,
5 µg plasmid, 30 units T7 RNA polymerase (Promega
Biotec, Madison, WI) and diethylpyrocarbonate (DEPC)-
treated water to a final volume of 100 µl. T7 RNA
20 polymerase was used in the transcription of the plasmid
DNA, because a T7 promotor, rather than the SP6
promotor, was found ahead of the 5'-end of the ligated
AR-cDNA.

25 The reaction was allowed to proceed for 2 hrs.
at 40°C. RQ1 DNase I (5 units) was added and the
reaction continued for 15 mins. at 40°C. The reaction
mixture was extracted with phenol/chloroform (1:1) and
then with chloroform. RNA product was precipitated by
the addition of 0.1 volume of 3 M Na-acetate and 2.5
30 volumes of ethanol, re-suspended in 0.5 M NaCl, and re-
precipitated with 2.5 volumes of ethanol. RNA trans-
cribed was isolated and then translated in a rabbit
reticulocyte lysate system.

35 Translation of RNA was carried out in a micro-
coccal nuclease-treated rabbit reticulocyte lysate
(Promega Biotec, Madison, WI) pre-mixed kit (100 µl) in
the presence of 8 µg mRNA, 40 µCi of [³⁵S] methionine

- 18 -

(800 Ci/mmol; Amersham Co., Arlington Heights, IL) and 100 μ M each of amino acid mixture without methionine. The reaction was allowed to proceed for 1 hour at 30°C. To quantitate the incorporation of radioactive methionine, 3 μ l of the reaction mixture were added to 1 ml of 1 M NaOH containing 1.5% H₂O₂, 1 mM methionine, and 0.04% bovine serum albumin. The mixture was incubated for 15 mins. at 37°C to hydrolyze [³⁵S] methionine charged tRNA. The radioactive protein products were precipitated by the addition of 1 ml of 25% trichloroacetic acid and the radioactivity associated with the precipitates was determined.

By SDS-PAGE (8% acrylamide gel) analysis, performed as described in Saltzman, et al., J. Biol. Chem., 262:432 (1987), it was found that a 79 kD protein comprised more than 85% of the translated products.

EXAMPLE 5

Binding Activity of the 79 kD hAR Protein to a Synthetic Androgen

To study the steroid binding activity of the protein coded for by the cloned cDNA, the reticulocyte lysate of Example 4, containing the newly synthesized protein was incubated with 17 α -[³H]-methyl-17 β -hydroxy-estra-4,9,11-trien-3-one ([³H] R1881), a potent synthetic androgen that binds AR with high affinity [Liao, et al., J. Biol. Chem., 248:6154 (1973)].

Specifically, RNA transcribed from the cloned cDNA, as described in Example 4, was translated in a rabbit reticulocyte lysate system and aliquots of the lysate were then incubated with 5 nM [³H] R1881 (87 Ci/mmol) in the absence or presence of 25 nM, 50 nM, or 250 nM of non-radioactive steroid. The final incubation volume was 100 μ l. The radioactive androgen binding was measured by the hydroxylapatite-filter method as des-

- 19 -

cribed in Liao, S., et al., J. Steroid Biochem., 20:11 (1984). The result was expressed as a percentage of the radioactivity bound in the control tube (5000 dpm) without additional non-radioactive steroid and is listed in Table 1.

TABLE 1
Androgen-specific binding of
hAR coded by cloned cDNA

Non-radioactive steroid added	[³ H] R1881-bound (% of control)		
	25 nM	50 nM	250 nM
R1881	13	10	1
5 α -dihydrotestosterone	25	17	6
5 β -dihydrotestosterone	89	89	81
17 β -Estradiol	91	91	86
Progesterone	100	91	92
Dexamethasone	100	93	93
Hydrocortisone	96	90	90
Testosterone	38	28	Not tested

As shown in Table 1, the active natural androgen, 17 β -hydroxy-5 α -androstane-3-one (5 α -dihydrotestosterone) competed well with [³H] R1881 binding, but the inactive 5 β -isomer did not compete well with [³H] R1881 suggesting that it does not bind tightly to AR. The binding activity was steroid specific; dexamethasone, hydrocortisone, progesterone, and 17 β -estradiol did not compete well with the radioactive androgen for binding to the 79 kD protein.

Similar steroid binding specificities have also been observed for rAR coded for by cloned cDNA. Chang, C., et al., Proc. Nat'l. Acad. Sci. (USA), 85:7211-7215 (1988).

Using the hydroxylapatite filter assay method, it was observed that approximately one molecule of the

- 20 -

35S-labelled 79 kD protein obtained from the lysate bound about one molecule of the tritiated androgen at a saturating concentration of ligand. By Scatchard plot analysis, the apparent dissociation constant was 0.31 nM, which is similar to the binding constant (0.65 nM) reported previously for AR of rat ventral prostate as reported in Schilling, et al., The Prostate, 5:581 (1984).

10

EXAMPLE 6Binding Activity of the 79 kD Protein to Human Auto-antibodies

It has previously been reported [Liao, et al., Proc. Nat'l. Acad. Sci. (USA), 82:8345 (1985)] that some older men with prostate cancers have high titers of auto-immune antibodies to AR in their serum samples. The ability of human auto-antibodies to recognize the 79 kD protein made by the reticulocyte lysate system was therefore studied. The receptor protein made in the lysate system of Example 4 was incubated with [³H] R1881 to allow the formation of radioactive androgen-androgen receptor (A-AR) complexes and was then mixed with serum containing auto-antibodies.

Reticulocyte lysate containing translated AR was incubated with [³H] R1881, as described in Example 4, and then incubated again in either the presence of or absence of 5 µl of human male serum containing antibodies to AR (anti-AR serum) for 4 hrs. at 4°C. Rabbit serum containing anti-human immunoglobulins (Anti-IgG) was then added as the second antibody. After 18 hrs. of incubation at 4°C, the mixture was centrifuged and the radioactivity associated with the precipitate was estimated. Human female serum, not containing anti-AR antibody, was also used for comparison.

- 21 -

The results shown in Table 2 below, indicate a quantitative immunoprecipitation of the radioactive A-AR complexes in the presence of both the high titer human serum and a rabbit anti-human immunoglobulin IgG. By SDS-PAGE, it was also observed that the immunoprecipitated protein was the 79 kD protein.

TABLE 2

10	Anti-human immunoglobulin-dependent precipitation of hAR made by the translation of RNA transcribed from cloned cDNA		
15	Sample incubated with [³ H]R1881	Anti-serum addition	Immunoprecipitable radioactivity(dpm)
	AR coded by cDNA ^a	None	32
		+Anti-AR serum + Anti-IgG	8212
		+Female serum + Anti-IgG	430
		+Anti-IgG	8
20		+Anti-AR serum + Anti-IgG	42
	Heated AR ^b BMW-lysate ^c	+Anti-AR serum + Anti-IgG	204

^a 8500 dpm of the radioactive AR complexes made were used.

^b Reticulocyte lysate containing AR was heated at 50°C for 20 mins. to inactivate receptor and release the radioactive androgen bound before the addition of antiserum.

^c Brome Mosaic Virus RNA was used in the reticulocyte lysate translation system instead of RNA transcribed from cloned cDNA.

- 22 -

EXAMPLE 7Characterization of TR2-cDNA

Of the more than 40 TR2-type human cDNA clones
5 obtained, including the 30 described in Example 1, the
clone designated TR2-5 was found to be 2029 base pairs
in length as indicated in Figure 4. The open reading
frame between the first ATG and terminator TAA can
encode 483 amino acids with a calculated molecular
10 weight of 52 kD. The putative DNA binding region is
underscored. The putative initiator ATG matched closely
with Kozak's consensus sequence for active start
codons. [See, Kozak, M., Nature, 308:241 (1984).] Two
triplets upstream of this ATG codon is an in-frame
15 terminator (TAA) further supporting initiator function
for the ATG. Eleven out of the 30 TR2-type clones of
Example 1, as represented by the clone designated TR2-7,
contain an internal 429 bp insertion between nucleotide
sequence 669 and 670 (designated by an asterisk in
20 Figure 4). This internal insertion introduces a
termination codon TAG (underscored in the insert
sequence footnote) which reduces the open reading frame
to 184 amino acids with a calculated molecular weight of
20 kD. It is likely that the insertion in these 11 TR2
25 clones (or deletion in the 19 other TR2 clones)
represents either the existence of two types of mRNA in
the human testis or an artifact of cDNA construction.
In the 3'-nontranslated region, a eukarotic
polyadenylation signal AATAAA is present between the
30 nucleotide sequence 2000 and 2007 of the TR2-5 clone.

Other variants of TR-2 with open reading
frames at the putative ligand-binding domains have been
obtained. Some of these may code for receptors for new
hormones or cellular effectors. It is anticipated that
35 the knowledge of TR2-cDNA sequences will be utilized in
isolation and structural analysis of other cellular

- 23 -

receptors, their genes, and ligands (endogenous or therapeutic agents) that can regulate cellular growth and functions in both normal and diseased organs.

Figure 5 depicts an amino acid sequence alignment of the cysteine-rich DNA binding domain of human
5 androgen receptor, glucocorticoid receptor, mineralocorticoid receptor, progesterone receptor, estrogen receptor, human TR2 protein, rat AR, chick vitamin D receptor (c-VDR), and the v-erb A oncogene product of avian
10 erythroblastosis virus. The numbers in the left margin represent the positions of amino acid residues in the individual receptors. Common residues are boxed with solid lines. The residues in dotted boxed represent those not in common with those in the solid boxes.
15 V-erb A has two more amino acids at the starred position.

In this region, the human and rat cDNAs for AR have identical amino acid sequences, although for some amino acids different codons are employed. Also in this
20 region, the homology between human AR or rat AR and other receptors is as follows: glucocorticoid receptor (GR), 76.4%; mineralo-corticoid receptors (MR), 76.4%; progesterone receptors (PR), 79.2%; estrogen receptors (ER), 55.6%; TR2, 45.8%; chick vitamin D receptor (c-VDR), 40.3%; and the v-erb A oncogene product of avian
25 erythroblastosis virus, 40.3%. In the putative region for steroid binding, which has about 200 amino acids near the -COOH terminal of steroid receptors, the homology between human AR or rat AR and hGR, hMR, or hPR
30 is about 45-55%, whereas the homology between human AR and rat AR and hER is less than 20%. Thus, human and rat AR appear to be more closely related to GR, MR, and PR, than to v-erb A or to receptors for estrogen, vitamin D, and thyroid hormones.

35 The DNA binding domain of TR2 (amino acids 111 to 183) has a high homology with the steroid receptor

- 24 -

super-family as follows: retinoic acid receptor (RAR), [Giguere, et al., Nature, 330:624 (1987)], 65%; thyroid receptor (T₃R) [Sap, et al., Nature, 324:635 (1987)], 59%; mineralocorticoid receptor (MR), [Arriza, et al., Science, 235:268 (1987)], 54%; vitamin D₃ receptor (VD₃R) [McDonnell, et al., Science, 235:1214 (1987)], 53%; hERR1 and hERR2, [Giguere, V., et al., Nature, 331:91 (1988)], 51% estrogen receptor (ER), [Hollenberg, et al., Nature, 318:635 (1985)], 51%; glucocorticoid receptor (GR) [Hollenberg, et al., Nature, 318:635 (1985)], 50%; androgen receptor (AR), 50%; progesterone receptor (PR), 49%; [Loosfelt, et al., Proc. Nat'l. Acad. Sci., (USA), 83:9045 (1986)]. As noted in Figure 5, the positions of 20 amino acids (9 Cys, 3 Arg, 2 Gly, 2 Phe, 1 Lys, 1 Met, 1 Asp, 1 His) in the putative DNA binding domain are identical among all isolated thyroid steroid receptor genes. It has been proposed that this highly conserved region may be involved in the formation of a DNA binding finger. See, Weinberger, et al., Nature, 318:670 (1985). Like the other steroid receptors, TR2 does not have the two extra amino acids (Lys-Asn) found only in the thyroid receptors' DNA binding domain. See, Sap, et al., Nature, 324:635 (1987).

EXAMPLE 8

In Vitro Transcription and Translation of TR2 cDNA

The Eco RI-Eco RI DNA inserts from clones TR2-5 and TR2-7 were isolated and ligated to an EcoRI digested pGEM-3Z vector for in vitro transcription essentially as described in Example 3. E. coli DH5 α cells, transformed with these plasmids were designated EC TR2-5 and EC TR2-7 and were deposited with the American Type Culture Collection, 12301 Parklawn Drive,

- 25 -

Rockville, Maryland 20852 on January 25, 1989 under Accession Nos. 67877 and 67876.

Transcribed RNA was then translated in a rabbit reticulocyte lysate system. By SDS-
5 polyacrylamide gel electrophoresis (PAGE), it was found that the major translated product of TR2-7, which has an internal 429 bp, insertion, was a 20 kD protein. The major translated product of TR2-5 was a 52 kD protein.

To further characterize these translated pro-
10 teins, the translation lysate was passed over a DNA cellulose column. The bound product was then eluted, concentrated and applied to SDS-PAGE. The results indicated that the translated proteins were indeed DNA-binding proteins.

15

EXAMPLE 9

Binding Activity of TR2-5 cDNA Expression Product

20 To study the steroid binding activity of the translation products of the TR2-5 clone, the products were incubated with all major classes of steroids, including androgens, progesterone, glucocorticoid and estrogen but no significant binding with the above
25 steroids was observed. This does not necessarily rule out a steroid binding function for the protein. Possibly the TR2-5 expression product steroid binding activity may involve some post-translation modifications missing in the rabbit reticulocyte lysate system.
30 Alternatively, the TR2-5 translated protein may be steroid independent or may bind to an unidentified ligand present in the human testis or rat ventral prostate.

The size of TR2 mRNA was determined by
35 Northern blot analysis with TR2-5 cDNA insert as a probe. One 2.5 kb band was detected which should

- 26 -

include enough sequence information to code for a 52 kD protein. The TR2 mRNA tissue distribution was also analyzed by dot hybridization. The hybridization was visualized by densitometric scanning of the autoradiographs, individual dots were cut and radioactivity measured by liquid scintillation counting [See, Chang, et al., J. Biol. Chem., 262:2826 (1987)]. The results showed that TR2 mRNA was most abundant in the rat ventral prostate with the relative amounts in other tissues being: prostate 100%, seminal vesicle 92%; testis, 42%; submaxillary gland, 18%; liver, 13%; kidney, <1%; and uterus, <1%.

EXAMPLE 10

Analysis of Androgen Regulation of AR and TR2 mRNA Levels in the Rat Ventral Prostate

Because rat ventral prostate is an androgen-sensitive organ and contains the greatest amount of AR and TR2 mRNA, the effect of androgen depletion and replacement on the mRNA levels was studied by RNA dot hybridization and Northern blot analysis. Total RNA was extracted from the ventral prostate of normal rats, rats castrated and rats previously castrated and treated with 5 α -dihydrotestosterone (17 β -hydroxy-5 α -androstand-3-one). AR mRNA levels per unit of DNA increased 200 to 300% of the level for normal rats within 2 days after castration. Administration of 5 α -dihydrotestosterone (5 mg/rat/day) into castrated rats reduced the AR mRNA level to that of normal rats. TR2 mRNA levels, per unit of DNA, were increased to 170% of the normal rat within 2 days after castration. Injection of 5 α -dihydrotestosterone (5 mg/rat/day) into castrated rats reduced the TR2 mRNA to the levels of normal rats. Interestingly, the total prostate RNA levels, at the same period of time, were decreased to 40% of the normal level. The

- 27 -

effects of androgen on the levels of prostatic TR2 mRNA were further confirmed by flutamide injection experiments. Flutamide, an anti-androgen which antagonizes the effects of 5 α -dihydrotestosterone on the ventral prostate weights in castrated rats [Neri, et al., Invest. Urol., 10:123 (1972)], was injected into normal rats for from 2 to 6 days. TR2 mRNA levels were then measured by dot hybridization as described above. The results show that flutamide injection, like castration, increased TR2 mRNA levels. The change in the AR or TR2 protein levels could be due to a change in mRNA stability and utilization or a change in the regulation of gene transcription. The activation or inactivation by androgen of specific genes to different degrees in the same organ may suggest that androgen is involved in the structuring of the pattern of gene expression in the target cell. Also, if androgen-mediated gene repression mechanisms are related to growth of the prostate, then a further study of the mechanism and structure of genes, repressed AR and TR2 mRNA may provide a better understanding of androgen action in the normal and abnormal prostate and other hormone sensitive organs.

Also, defects in the structures of AR and androgen sensitive genes and/or losses of the control of the production and function of these gene products can be the causes of the abnormal growth of androgen sensitive or insensitive tumors like prostate cancers. These lines of research may, therefore, be helpful in designing new diagnostic methods and treatments for patients.

- 28 -

EXAMPLE 11Expression of Cloned AR-Genes
and Androgen Sensitive Genes
in Eukaryotic and Prokaryotic Cells

5 The ability of cloned genes to function when
introduced into mammalian, yeast, and bacterial cells
has proved to be very valuable in understanding the
function and regulatory mechanism of genes. Recombinant
10 techniques can provide, in large quantities, gene
expression products (proteins) which are not readily
obtainable from natural sources. While bacterial
systems are very useful in large scale production of
those proteins which do not require substantial post-
15 translational modification for optimal biological
activity, eukaryotic systems are particularly
advantageous because of their ability to correctly
modify the expressed proteins to their functional forms.

 Using well known techniques, AR-cDNA and TR2-
cDNA may readily be used for large scale production of
20 gene products. For this purpose, the most efficient
transcription units can be constructed using viral, as
well as non-viral, vectors with regulatory signals that
can function in a variety of host cells. SV40, pSV2,
adenoviruses, and bovine papilloma virus DNA have been
25 used successfully for introduction of many eukaryotic
genes into eukaryotic cells and permit their expression
in a controlled genetic environment. These and similar
systems are expected to be appropriate for the
expression of AR- and TR2-genes. To assist gene
30 transfer, the two most widely used methods, the "calcium
phosphate precipitation" and the "DEAE-dextran
technique" can be used. Genes can be introduced into
cells either transiently, where they continue to express
for up to 3 days, or, more permanently to form stably
35 transformed cell-lines. The expressed proteins can be
detected by androgen binding or antibody assays.

- 29 -

The expression of cloned AR-genes was achieved as follows in a eukaryotic system. NIH 3T3 cells, contact-inhibited cells established from NIH Swiss mouse embryo, were co-transfected with hAR cDNA inserted into pBPVMTH vectors as described by Gorman, "DNA Cloning", 2:143-190 D. M. Glover, ed.; (Oxford, Washington, D.C. 1985). Transfected cells were cloned and grown in multiple-well cell culture plates. About 100 individual cell lines were isolated. Of these, 6 demonstrated [³H] R1881-binding activity at least 4-fold the activity of cells transfected with pSV2 vector alone, i.e., without the hAR cDNA sequence.

To express AR cDNA in prokaryotic systems, hAR and rAR cDNAs were inserted into a number of expression vectors including pUR, λ GT11, pKK223-3, pKK233-2, pLEX, pATH1, pATH2, pATH10, and pATH11. Vectors with AR cDNA inserts were used to infect E. coli strains (JM109, DH5 α , Y1089, JM105, and RR1). According to polyacrylamide gel electrophoresis analysis, the infected bacteria can synthesize AR fragments coded for by the AR cDNA inserts. Some of these AR polypeptides are degraded in culture. Amino terminal, DNA-binding, and androgen binding domains were used, as described in Example 12, to construct fusion proteins representing these domains.

EXAMPLE 12

Production of Polyclonal and Monoclonal Antibodies to AR

The isolation of AR in significant amounts from androgen sensitive organs has been exceedingly difficult. Therefore, the high-level expression of hAR or rAR cDNAs, as shown in Example 11, is expected to be an ideal way for the large scale production of AR. In addition, oligopeptides, with sequences identical to the

- 30 -

deduced amino acid sequences of portions of AR molecules, can be chemically synthesized inexpensively in large quantities. Both AR produced by expression vectors in eukaryotic or prokaryotic cells and AR oligopeptides chemically synthesized were used as antigens for the production of monoclonal antibodies as described in greater detail below.

Generally, several chemically synthesized oligopeptides, representing sequences unique to AR, (i.e., PYGDMRLETARDEHVLV; CPYGDMRLETARDHVLP; and SIRRNLVYSCRGSKDCIINK) were bound to BSA or KLH carrier proteins and were used to immunize mice. Spleen cells from these mice were fused to myeloma cells to produce hybrid antibody producing cells. Analysis by ELISA (enzyme-linked immunoassay) of the supernatants of 4 hybrid cultures appeared to indicate the presence of immunoglobulin that interacts with AR of rat ventral prostate. It is anticipated that these cells which produce monoclonal antibodies can be injected intraperitoneally into BALB/c mice previously treated with pristane. Ascites fluids can then be harvested and antibodies precipitated with ammonium sulfate.

25 Expression of Androgen Receptor Fusion Protein in E. coli

Three different parts of the AR gene (encompassing the N-terminal domain, the DNA-binding domain and the androgen-binding domain) were fused, in frame, to the N-terminal half of the trpE gene (trpE promoter-the first 969 bp of trpE coding region-multiple cloning region of pUC12) by using the PATH expression vectors as shown in Figures 6, 7, and 8, respectively. Dieckmann, et al., J. Biol. Chem., 260:1513 (1985).

These constructions resulted in the fusion of approximately 25 kDa of AR, including a portion of the N-terminal domain; 29 kDa of AR, including a major

- 31 -

portion of the DNA-binding domain; and 12 kDa of AR, including a portion of the androgen-binding domain; to the 33 kDa trpE protein. Because the trpE protein is insoluble, partially purified induced fusion proteins were obtained simply by lysing the E. coli and precipitating the insoluble fusion proteins. After electrophoresis on SDS-polyacrylamide gels, the induced fusion proteins, i.e., those proteins not present in the control pATH vector (no AR gene insert), were sliced from the gels and then used for immunization.

Fusion proteins, other than the three specifically exemplified, can also be constructed using these means.

15 Production and Purification of Anti-AR Antibodies

Rabbits, rats, and mice were immunized with either SDS-polyacrylamide gel slices containing denatured fusion proteins or electro-eluted, SDS-free, fusion protein, as well as fusion proteins obtained by other protein purification methods. The presence of antibodies to the fusion proteins in the antisera was assayed by ELISA. Positive serum having a higher titer was further assayed by the double antibody precipitation method using rat ventral prostate cytosol [³H]AR as antigen. The results showed that 1 μ l of crude serum precipitated 10 to 20 fmole [³H]AR. Anti-AR crude serum was then affinity-purified by differential suspension of immune serum containing TrpE protein(s) (both those TrpE proteins having and those TrpE proteins not having inserted AR sequences) expressed by pATH vectors. The bound antibodies can be removed from the suspension because TrpE protein is insoluble. Antibodies specific against only the trpE protein were removed; antibodies specific for AR were isolated and again confirmed by both ELISA and double antibody precipitation.

- 32 -

Production of Monoclonal
Anti-Androgen Receptor Antibodies

5 The immunized rats were judged ready to be
sacrificed for a fusion when their serum tested positive
anti-AR antibodies by ELISA. Spleens were removed and
grinded to release the cells into DMEM (Dulbeco's
Modified Engle's Medium) medium. Through a series of
centrifugations using DMEM + DMEM with Ficoll Hypaque,
10 the spleen cells were isolated. The SP2/0 myeloma cells
were grown, split and diluted in 50 ml of DMEM with 20%
FCS, 1% MOPS, and 1X L-Gln for two days before ready for
the fusion. SP2/0 cells (5×10^6) and 5×10^7 spleen
cells were used in the fusion. After incubating
15 overnight, the fused cells were collected, suspended in
DMEM with 1X H-T, 1X Methotrexate, 20% FCS, and 1X PBS
and distributed in 96-well plates. Plates were
supplemented after 6 days with DMEM and 20% FCS.
Hybridomas were identified and assayed, using the ELISA
20 assay of Engrall, et al., Bio. Chem. et Biophys. ACTA,
251:427-439 (1971). In this assay, plates were coated
with either the AR fusion proteins or the TrpE protein
as antigen and read on an ELISA reader.

25 Only those hybridomas that caused a positive
reaction with the AR fusion protein were "limit diluted"
to a concentration of 10 cells/ml and were then
distributed among half of a 96-well plate. The
remaining cells from the original well were transferred
to a 24-well plate. Each of these plates had a
30 thymocyte feeder layer. The thymocyte feeder layer was
made up of thymus cells isolated from an un-injected
rat, purified through centrifugation, irradiated with
1200 to 1400 RADS, and diluted to 1×10^7 cells/ml of
DMEM with 20% FCS.

35 Positives from these thymocyte 96-well plates
were again tested by ELISA. Only those which again

- 33 -

tested positive with the AR fusion protein were grown up for monoclonal antibody purification. Three of the wells produced monoclonal antibody against AR. Both ELISA and double antibody assays were positive. The monoclonal antibodies were designated AN1-6, AN1-7, and AN1-15 and the three cell lines were designated HAN1-6, HAN1-7, and HAN1-15; Accession Nos. 10,000; 9,999; and 10,001; respectively, deposited on January 25, 1989 with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852.

Specificity of Anti-AR Antibodies

Sucrose gradient centrifugation was used to characterize the specificity of the three monoclonal anti-AR antibodies and their ability to react with non-denatured [^3H]AR.

Cytosol was prepared from the ventral prostates of castrated rates as follows. Rats were castrated by the scrotal route while under anesthesia. They were killed 18 hrs. later by cervical dislocation and their ventral prostates were removed, minced with scissors, washed in Buffer A (50 mM sodium phosphate, pH 7.5, 1 mM EDTA, 2 mM DTT, 10 mM sodium molybdate, 10% (v/v) glycerol and 10 mM sodium fluoride) and homogenized in 2x the tissue volume of Buffer A + 0.1 mM bacitracin, 1 mM PMSF, and aprotinin (1TIU/ml). The homogenate was centrifuged at 5,000 x g for 10 mins., adjusted to 10 nM ^3H -androgen, spun at 225,000 x g for 45 mins. and treated with dextran-coated charcoal. One hundred μl of the cytosol solution, containing ^3H -A-AR complexes, was incubated for 6 hrs. with 100 μl of the purified anti-androgen receptor monoclonal antibody, AN1-6, (20x as concentrated as the tissue culture media). Sucrose gradient centrifugation was performed by centrifugation at 257,000 x g for 16 hrs. at 4°C on a 3.8 ml, linear

- 34 -

5-20% (w/v) sucrose gradient containing 20 mM Tris-HCl, pH 7.5, 1 mM EDTA, 1 mM DTT, 10% (v/v) glycerol, and 0.4 M KCl. Gradients were fractionated and numbered from the bottom and 0.2 ml per fraction collected. The results obtained indicated that all three of the monoclonal antibodies, AN1-6, AN1-7, and AN1-15, recognized and effectively bound the radioactively labeled androgen receptor ([³H] AR).

The [³H]AR and other steroid receptor complexes had a sedimentation coefficient of about 4-5S in the sucrose gradient media containing 0.4M KCl. Anti-AR antibodies do not alter the sedimentation coefficient of 4-5S for [³H]glucocorticoid receptors complexes of rat liver, estrogen receptor complexes of MCF-7 cells, and progesterone receptor complexes of T47D cells, but do shift the sedimentation coefficient of [³H]A-AR complexes of rat ventral prostate from 4S to 9-12S or to heavier units. By SDS-polyacrylamide gel electrophoresis analysis it was also found that all major in vitro transcription/translation products of human and rat AR cDNAs were immunoprecipitable by the anti-AR antibodies.

EXAMPLE 13

Use of AR cDNA and TR2 cDNA as Probes
in the Study of Abnormality in Human
and Animal Organs and Cancer Cells

Patients with metastatic prostatic cancer initially often respond favorably to androgen withdrawal therapy (castration or antiandrogen treatments). Most patients, however, eventually relapse to an androgen-state for which no chemotherapy, which would significantly increase the survival rate, is available. Regardless of the origin of androgen-independent or -insensitive cancer cells, it is important to understand whether the androgen

- 35 -

insensitivity or abnormality in the diseased cells are due to qualitative or quantitative changes in (a) the AR or TR2 genes, (b) regulation of their transcription, or translation, or (c) other cellular factors. AR cDNA, TR2 cDNA, or their partial segments can be used as specific probes in these studies.

For the analysis of AR or TR2 genes, high molecular weight genomic DNA isolated from target organs, tumors, and cultured cells can be used in identifying and characterizing AR genes. Different restriction endonucleases can be used to cleave DNA. The fragments can be analyzed by Southern analysis (agarose electrophoresis, transfer to nitrocellulose and hybridization with AR cDNA probes). After identification, selected fragments can be cloned and sequenced. It is also possible to use appropriate oligonucleotide fragments of AR or TR2 cDNA as primers to amplify genomic DNA isolated from normal and abnormal organs or cells by specific DNA polymerases. The amplified genomic DNA can then be analyzed to identify sequence abnormality using the polymerase chain reaction (PCR) assay. Saiki, et al., Science, 230, 1350 (1985). See also, Mullis, K.B., U.S. Patent No. 4,683,202; July 28, 1987; and Mullis, K.B., U.S. Patent No. 4,683,195; July 28, 1987. For the analysis of mRNA for ARs or related proteins, dot hybridization and Northern hybridization analysis could be used to characterize mRNA and AR or receptor-like molecules quantitatively and qualitatively. From these studies valuable information about the number of different forms of AR genes and their expression in androgen insensitive and sensitive tumor cells can be obtained.

DNAs and RNAs obtained from androgen sensitive and insensitive tumors and from cell lines from rats and humans with testicular feminization syndromes have been analyzed by the above methods. Preliminary studies

- 36 -

indicated that abnormality in androgen responses may be due to sequence deletion/mutation in genes for ARs.

EXAMPLE 14

5

Development of Transgenic Animals

Transgenic techniques have been employed for expression of exogenous DNA. It may therefore be possible to confer androgen sensitivity to animals with androgen receptor defects. For example, androgen insensitive animals, such as testicular feminized mice or rats, are known to have defective AR genes or defective AR itself. If DNA containing a normal AR gene is injected into fertilized mouse embryos, the transgenic mice may carry and express the gene and produce a functional AR necessary for androgen responses. For micro-injection, it is necessary to use AR genes containing DNA that can be expressed in the insensitive animals.

20 A number of genomic receptor clones from human X-chromosome libraries and rat genomic DNA libraries have been obtained and analyzed for their structures. Clones containing AR sequences will be characterized by endonuclease mapping, by Southern hybridization and by
25 S1-nuclease mapping. The 5' and 3' untranslated regions thus identified will aid in determining the minimal size of the DNA that would be required for tissue specific expression of the AR coding region.

Partial sequence analysis of the 5' and 3' regions would locate the minimal region that represents the promoter and the polyadenylation region. Approximately 2 to 5 kb of upstream un-translated region and 0.5 to 1 kb of sequences downstream from the poly(A) site may be fused to the cDNA clone (minimal-gene) and
35 injected into embryos of mice. Transgenic mice would be identified by analysis of their tail DNA using mini-gene specific probe(s).

- 37 -

Normally only some of the transgenic mouse lines can express their transgenes. Transgenes may be inactive because of the presence of inhibitory sequences, integration of the exogenous gene into a transcriptionally inactive chromosomal location, or the juxtaposition of the transgene and an endogenous enhancer. In addition, androgen insensitivity may be due to various other factors and not due to abnormality in the AR gene or its expression.

The foregoing illustrative examples relate to the isolation of human and rat cDNAs encoding DNA binding proteins including androgen receptor and TR-2 and more particularly describe the transcription of the corresponding cDNAs and translation of the corresponding mRNAs in cell-free systems. While the present invention has been described in terms of specific methods and compositions, it is understood that variations and modifications will occur to those skilled in the art upon consideration of the present invention.

Accordingly it is intended in the appended claims to cover all such equivalent variations which come within the scope of the invention as claimed.

25

30

35

- 38 -

WHAT IS CLAIMED IS

1. A purified and isolated DNA sequence encoding androgen receptor polypeptide.
5
2. The DNA sequence according to claim 1 encoding human androgen receptor polypeptide.
3. The DNA sequence according to claim 1 encoding rat androgen receptor polypeptide.
10
4. A purified and isolated DNA sequence encoding TR2 polypeptide.
5. The DNA sequence according to claim 1 or 4 which is a cDNA sequence.
15
6. The DNA sequence according to claim 1 or 4 which is a genomic DNA sequence.
20
7. The DNA sequence according to claim 1 or 4 which is a partially synthetic DNA sequence.
8. The DNA sequence according to claim 1 and as set forth in Figure 3.
25
9. The DNA sequence according to claim 4 and as set forth in Figure 4.
10. A procaryotic or eucaryotic host cell transformed or transfected with a DNA sequence according to claim 1 or 4.
30

35

- 39 -

11. The procaryotic transformed host cell according to claim 10 which is E. coli DH5a cells designated as, and corresponding to A.T.C.C. deposit Nos.: EC-hAR 3600, A.T.C.C. No. 67879; EC-rAR 2830, 5 A.T.C.C. No. 67878; EC TR2-5, A.T.C.C. 67877; and EC TR2-7, A.T.C.C. No. 67876.

12. A viral or circular DNA plasmid comprising a DNA sequence according to claim 1 or 4.

10

13. A viral or circular DNA plasmid according to claim 11 further comprising an expression control DNA sequence operatively associated with said androgen receptor or TR2 encoding DNA.

15

14. A method for the production of androgen receptor polypeptide comprising:

growing, in culture, a host cell transformed or transfected with a DNA sequence according to claim 1; 20 and

isolating from said host cell or culture the polypeptide product of the expression of said DNA sequence.

25 15. A method for the production of androgen receptor polypeptide comprising:

disposing a DNA sequence according to claim 1 in a cell free transcription and translation system; and isolating from said system the polypeptide 30 product of the expression of said DNA sequence.

16. A method for the production of TR2 polypeptide comprising:

growing, in culture, a host cell transformed or transfected with a DNA sequence according to claim 4; 35 and

- 40 -

isolating from said host cell or culture the polypeptide product of the expression of said DNA sequence.

5 17. A method for the production of TR2 polypeptide comprising:
 disposing a DNA sequence according to claim 4
 in a cell free transcription and translation system; and
 isolating from said system the polypeptide
10 product of the expression of said DNA sequence.

 18. The polypeptide product of the in vitro
 or in vivo expression of a DNA sequence according to
 claim 1.

15

 19. An amino acid sequence as set out in
 Figure 3.

 20. The polypeptide product of claim 18
20 characterized by a molecular weights of 98 kD and 79 kD
 by SDS-PAGE and the ability to bind an androgen.

 21. The polypeptide product of the in vitro
 or in vivo expression of a DNA sequence according to
25 claim 4.

 22. TR2 polypeptides.

 23. A synthetic peptide duplicative of a
30 sequence of amino acids present in AR or TR2 proteins in
 a region of the proteins not involved with DNA binding
 functions and sharing at least one antigenic epitope
 with AR or TR2 proteins.

35

- 41 -

24. An antibody specifically immunoreactive with at least one epitope of androgen receptor polypeptide or TR2 polypeptide other than an epitope within the DNA binding functional region thereof.

5

25. The monoclonal antibody according to claim 24.

26. The monoclonal antibody according to claim 24 and produced by hybridoma cell line Nos. HB 10,000; HB 9,999; and HB 10,001.

10

27. The polyclonal antibody according to claim 24.

15

28. A method for quantitative detection of androgen receptor based on the immunological reaction of androgen receptor with an antibody according to claim 24.

20

29. A method for quantitative detection of TR2 receptor based on the immunological reaction of TR2 receptor with an antibody according to claim 24.

25

30. A method for the quantitative detection of androgen receptor encoding DNA or RNA based on hybridization of said nucleic acids with a DNA sequence according to claim 1.

30

31. A method for the quantitative detection of TR2 receptor encoding DNA or RNA based on hybridization of said nucleic acids with a DNA sequence according to claim 4.

35

- 42 -

32. A method for the quantitative and qualitative detection of AR or TR2 specific gene sequence or sequences present in a sample comprising the steps of:

5 a) treating said sample with one oligonucleotide primer for each strand for said specific sequence, under hybridizing conditions such that for each strand of each sequence to which an oligonucleotide primer is hybridized an extension product of each primer
10 is synthesized which is complementary to each nucleic acid strand, wherein said primer or primers are selected so as to be sufficiently complementary to each strand of each specific sequence to hybridize therewith such that the extension product synthesized from one primer, when
15 it is separated from its complement, can serve as a template for synthesis of the extension product of the other primer;

 b) treating the sample under denaturing conditions to separate the primer extension products
20 from their templates if the sequence or sequences to be detected are present;


 c) treating the sample with oligonucleotide primers such that a primer extension product is synthesized using each of the single strands produced in
25 step (b) as a template, resulting in amplification of the specific nucleic acid sequence or sequences if present;

 d) adding to the product of step (c) a labeled oligonucleotide probe for each sequence being
30 detected capable of hybridizing to said sequence or a mutation thereof; and

 e) determining whether said hybridization has occurred.

35

1/14



SUBSTITUTE SHEET

2/14

FIGURE 2




FIGURE 5

h-GR 419	Lys	Leu	Cys	Leu	Val	Cys	Ser	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Val	Leu	Thr	Cys	Gly	Ser	Cys	Lys
h-MR 601	Lys	Ile	Cys	Leu	Val	Cys	Gly	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Val	Val	Thr	Cys	Gly	Ser	Cys	Lys
h-PR 565	Lys	Ile	Cys	Leu	Ile	Cys	Gly	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Val	Leu	Thr	Cys	Gly	Ser	Cys	Lys
h-AR	Lys	Thr	Cys	Leu	Ile	Cys	Gly	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Ala	Leu	Thr	Cys	Gly	Ser	Cys	Lys
r-AR	Lys	Thr	Cys	Leu	Ile	Cys	Gly	Asp	Glu	Ala	Ser	Gly	Cys	His	Tyr	Gly	Ala	Leu	Thr	Cys	Gly	Ser	Cys	Lys
h-ER 183	Arg	Tyr	Cys	Ala	Val	Cys	Asn	Asp	Tyr	Ala	Ser	Gly	Tyr	His	Tyr	Gly	Val	Trp	Ser	Cys	Glu	Gly	Cys	Lys
h-TR2	Asp	Leu	Cys	Val	Val	Cys	Gly	Asp	Lys	Ala	Ser	Gly	Arg	His	Tyr	Gly	Ala	Val	Thr	Cys	Glu	Gly	Cys	Lys
v-erbA35	Glu	Gln	Cys	Val	Val	Cys	Gly	Asp	Lys	Ala	Thr	Gly	Tyr	His	Tyr	Arg	Cys	Ile	Thr	Cys	Glu	Gly	Cys	Lys
c-VDR	Arg	Ile	Cys	Gly	Val	Cys	Gly	Asp	Arg	Ala	Thr	Gly	Phe	His	Asn	Ala	Met	Thr	Cys	Glu	Gly	Cys	Lys	Lys
h-GR 439	Val	Phe	Phe	Lys	Arg	Ala	Val	Glu	Gly	Gln	His	Asn	Tyr	Leu	Cys	Ala	Gly	Arg	Asn	Asp	Cys	Ile	Ile	Asp
h-MR 625	Val	Phe	Phe	Lys	Arg	Ala	Val	Glu	Gly	Gln	His	Asn	Tyr	Leu	Cys	Ala	Gly	Arg	Asn	Asp	Cys	Ile	Ile	Asp
h-PR 589	Val	Phe	Phe	Lys	Arg	Ala	Met	Glu	Gly	Gln	His	Asn	Tyr	Leu	Cys	Ala	Gly	Arg	Asn	Asp	Cys	Ile	Ile	Asp
h-AR	Val	Phe	Phe	Lys	Arg	Ala	Ala	Glu	Gly	Lys	Gln	Lys	Tyr	Leu	Cys	Ala	Ser	Arg	Asn	Asp	Cys	Thr	Ile	Asp
r-AR	Val	Phe	Phe	Lys	Arg	Ala	Ala	Glu	Gly	Lys	Gln	Lys	Tyr	Leu	Cys	Ala	Ser	Arg	Asn	Asp	Cys	Thr	Ile	Asp
h-ER 207	Ala	Phe	Phe	Lys	Arg	Ala	Ala	Glu	Gly	Lys	Gln	Lys	Tyr	Leu	Cys	Ala	Ser	Arg	Asn	Asp	Cys	Thr	Ile	Asp
h-TR2	Gly	Phe	Phe	Lys	Arg	Ser	Ile	Gln	Gly	His	Asn	Asp	Tyr	Met	Cys	Pro	Ala	Thr	Asn	Gln	Cys	Thr	Ile	Asp
v-erbA59	Ser	Phe	Phe	Lys	Arg	Ser	Ile	Arg	Lys	Asn	Leu	Val	Tyr	Ser	Cys	Arg	Gly	Ser	Lys	Asp	Cys	Val	Ile	Asp
c-VDR	Gly	Phe	Phe	Arg	Arg	Ser	Met	Lys	Arg	Lys	Ala	Met	Phe	Thr	Cys	Thr	Tyr	Asp	Gly	Cys	Cys	Val	Ile	Asp
h-GR 463	Lys	Ile	Arg	Arg	Lys	Asn	Cys	Pro	Ala	Cys	Arg	Tyr	Arg	Lys	Cys	Leu	Gln	Ala	Gly	Met	Asn	Leu	Glu	Ala
h-MR 649	Lys	Ile	Arg	Arg	Lys	Asn	Cys	Pro	Ala	Cys	Arg	Leu	Gln	Lys	Cys	Leu	Gln	Ala	Gly	Met	Asn	Leu	Gly	Ala
h-PR 613	Lys	Ile	Arg	Arg	Lys	Asn	Cys	Pro	Ala	Cys	Arg	Leu	Arg	Lys	Cys	Cys	Gln	Ala	Gly	Met	Val	Leu	Gly	Gly
h-AR	Lys	Phe	Arg	Arg	Lys	Asn	Cys	Pro	Ser	Cys	Arg	Leu	Arg	Lys	Cys	Tyr	Glu	Ala	Gly	Met	Thr	Leu	Gly	Ala
r-AR	Lys	Phe	Arg	Arg	Lys	Asn	Cys	Pro	Ser	Cys	Arg	Leu	Arg	Lys	Cys	Tyr	Glu	Ala	Gly	Met	Thr	Leu	Gly	Ala
h-ER 231	Lys	Asn	Arg	Arg	Lys	Ser	Cys	Gln	Ala	Cys	Arg	Leu	Arg	Lys	Cys	Tyr	Glu	Ala	Gly	Met	Thr	Leu	Gly	Ala
h-TR2	Lys	His	His	Arg	Asn	Arg	Cys	Gln	Tyr	Cys	Arg	Leu	Gln	Arg	Cys	Ile	Ala	Phe	Gly	Met	Met	Lys	Gly	Gly
v-erbA85	Lys	Ile	Thr	Arg	Asn	Gln	Cys	Gln	Leu	Cys	Arg	Phe	Lys	Lys	Cys	Cys	Ile	Ser	Val	Gly	Met	Ala	Met	Asp
c-VDR	Lys	Asp	Asn	Arg	Arg	His	Cys	Gln	Ala	Cys	Arg	Leu	Lys	Arg	Cys	Val	Asp	Ile	Gly	Met	Met	Met	Lys	Lys

SUBSTITUTE SHEET

10/14

[illegible]

CGC CCG GGG ATC CTC TAG
Arg Pro Gly Ile Leu STOP

5 amino acid linker

Total amino acid: $323 + 6 + 242 + 5 = 576$

FIGURE 6B

SUBSTITUTE SHEET

323 amino acids from TRP E protein

170 ATG CAA ACA CAA AAA CCG ACT CTC GAA CTC CTA ACC TGC GAA GGC CCT TAT CCG GAC AAT CCC ACC GCG CTT TTT CAC CAG TTG TGT GGG 230
180 MET GLN THR GLN LYS PRO THR LEU GLU LEU THR CYS GLU THR CYS GLY ALA LEU PHE HIS GLN LEU CYS GLY 240
250
260 GAT GGT CCG CCA ACG CTG CTG GCA GAT ATC GAC ACG AAA GAT CAT TTA AAA ACG CTG CTG GTA GAC AGT GCG CTG CCG 310
270 ASP ARG PRO ALA THR LEU LEU LEU SER ALA ASP ILE ASP SER LYS ASP LYS ASP LYS LEU LEU VAL ASP SER ALA LEU ARG 320
330
340
350 ATT ACA GCT TTA GGT GAC ACT CTC ACA ATC CAG GCA CTT TCC GCG AAC GCG GAA GCG CTC CTG GCA CTA CTG GAT AAC GCG CTG CCT GCG 410
360 ILE THR ALA LEU GLY ASP THR VAL THR ILE GLN ALA LEU SER GLY ASN GLY GLU ALA LEU LEU ALA LEU LEU ASP ASN ALA LEU PRO ALA 420
370
380
390
400
410
420
430
440 GGT GTG GAA GCT GAA CCA AAC TCC CCG GTG CCG TTC CCC CCT CTC ACT CCA CTG GAT GAA GAC GCG GCG TTA TGC TCC 490
450 GLY VAL GLU SER GLU GLN SER PRO ASN CYS ARG VAL LEU ARG PHE PRO PRO VAL SER PRO LEU LEU ASP GLU ASP ALA ARG LEU CYS SER 500
510
520
530 CTT TCG GTT TTT GAC GCT TTC CGT TTA TTG CAG AAT CTG TTG AAT GTA CCG AAG GAA GAA CCA GAA CCC ATG TTC ACC GCG CTG TTC 580
540 LEU SER VAL PHE ASP ALA PHE ARG LEU LEU LEU ASN LEU LEU VAL PRO LYS GLU ARG GLU ALA MET PHE PHE SER GLY LEU PHE 590
600
610
620 TCT TAT GAC CTT GTG CCG GGA TTT GAA GAT TTA CCG CAA CTG TCA CCG GAA AAT AAC TGC CCT GAT TTC TGT TAT CTC GCT GAA ACG 660
630 SER TYR ASP LEU VAL ALA GLY PHE LEU ASP LEU PRO GLN LEU SER ALA GLU ASN ASN CYS PRO ASP PHE CYS PHE TYR LEU ALA GLU THR 670
680
690
700
710 CTG ATG GTG ATT GAC CAT CAG AAA AAA ACG ACC CGT ATT CAG CCG ACC CTG TTT CCG AAT GAA GAA CAA AAA CAA CGT CTC ACT GCT 760
720 LEU MET VAL ILE ASP HIS GLN LYS LYS SER THR ARG ILE GLN ALA SER LEU PHE ALA PRO ASN GLU GLU LYS GLN ARG LEU THR ALA 770
780
790
800 CCG CTG AAC GAA CTA CCG CAG CAA CTG ACC GAA GCG CCG CCG CTG CCA CTG GTT TCC CTG CCG CAT ATG CGT TGT GAA TGT AAT CAG 840
810 ARG LEU ASN GLU LEU ARG GUN GUN LEU THR GLU ALA ALA PRO PRO LEU VAL SER VAL PRO HIS MET ARG CYS GLU CYS ASN GLN 850
860
870
880 ACC GAT GAA GAG TTC GGT GCG GTA GTG CGT TTG TTG CAA AAA CCG ATT CCG CCG GGA GAA ATT TTC CAG GTG CCA TCT CCG CCG TTC 910
890 SER ASP GLU GLU PHE GLY VAL VAL ARG LEU LEU GLN LYS ALA ILE ARG ALA GLY ILE PHE GLN VAL VAL PRO SER ARG ARG PHE 920
930
940
950
960
970
980 TCT CTG CCC TGC CCG TCA CCG CTG CCG GCG TAT TAC GTG CTG AAA AAG AGT AAT CCC ACC CCG TAC ATG TTT TTY ATG CAG GAT AAT GAT 1010
990 SER LEU PRO CYS PRO SER PRO LEU ALA ALA TYR TYR VAL, LEU LYS LYS SER ASN PRO SER PRO TYR MET PHE PHE MET GLN ASP ASP 1020
1030
1040
1050
1060
1070 TTC ACC CTA TTT CCG GCG TCG CCG GAA ACG TCG CTC AAG TAT GAT GCG ACC ACC CCG CAG ATT GAG ATC 1110
1080 PHE THR LEU PHE GLY ALA SER PRO GLU SER SER LEU LYS TYR ASP ALA THR SER ARG GLN ILE GLU ILE 1120
1130
1140
1150
1160
1170
1180
1190
1200
1210
1220
1230
1240
1250
1260
1270
1280
1290
1300
1310
1320
1330
1340
1350
1360
1370
1380
1390
1400
1410
1420
1430
1440
1450
1460
1470
1480
1490
1500
1510
1520
1530
1540
1550
1560
1570
1580
1590
1600
1610
1620
1630
1640
1650
1660
1670
1680
1690
1700
1710
1720
1730
1740
1750
1760
1770
1780
1790
1800
1810
1820
1830
1840
1850
1860
1870
1880
1890
1900
1910
1920
1930
1940
1950
1960
1970
1980
1990
2000
2010
2020
2030
2040
2050
2060
2070
2080
2090
2100
2110
2120
2130
2140
2150
2160
2170
2180
2190
2200
2210
2220
2230
2240
2250
2260
2270
2280
2290
2300
2310
2320
2330
2340
2350
2360
2370
2380
2390
2400
2410
2420
2430
2440
2450
2460
2470
2480
2490
2500
2510
2520
2530
2540
2550
2560
2570
2580
2590
2600
2610
2620
2630
2640
2650
2660
2670
2680
2690
2700
2710
2720
2730
2740
2750
2760
2770
2780
2790
2800
2810
2820
2830
2840
2850
2860
2870
2880
2890
2900
2910
2920
2930
2940
2950
2960
2970
2980
2990
3000
3010
3020
3030
3040
3050
3060
3070
3080
3090
3100
3110
3120
3130
3140
3150
3160
3170
3180
3190
3200
3210
3220
3230
3240
3250
3260
3270
3280
3290
3300
3310
3320
3330
3340
3350
3360
3370
3380
3390
3400
3410
3420
3430
3440
3450
3460
3470
3480
3490
3500
3510
3520
3530
3540
3550
3560
3570
3580
3590
3600
3610
3620
3630
3640
3650
3660
3670
3680
3690
3700
3710
3720
3730
3740
3750
3760
3770
3780
3790
3800
3810
3820
3830
3840
3850
3860
3870
3880
3890
3900
3910
3920
3930
3940
3950
3960
3970
3980
3990
4000
4010
4020
4030
4040
4050
4060
4070
4080
4090
4100
4110
4120
4130
4140
4150
4160
4170
4180
4190
4200
4210
4220
4230
4240
4250
4260
4270
4280
4290
4300
4310
4320
4330
4340
4350
4360
4370
4380
4390
4400
4410
4420
4430
4440
4450
4460
4470
4480
4490
4500
4510
4520
4530
4540
4550
4560
4570
4580
4590
4600
4610
4620
4630
4640
4650
4660
4670
4680
4690
4700
4710
4720
4730
4740
4750
4760
4770
4780
4790
4800
4810
4820
4830
4840
4850
4860
4870
4880
4890
4900
4910
4920
4930
4940
4950
4960
4970
4980
4990
5000
5010
5020
5030
5040
5050
5060
5070
5080
5090
5100
5110
5120
5130
5140
5150
5160
5170
5180
5190
5200
5210
5220
5230
5240
5250
5260
5270
5280
5290
5300
5310
5320
5330
5340
5350
5360
5370
5380
5390
5400
5410
5420
5430
5440
5450
5460
5470
5480
5490
5500
5510
5520
5530
5540
5550
5560
5570
5580
5590
5600
5610
5620
5630
5640
5650
5660
5670
5680
5690
5700
5710
5720
5730
5740
5750
5760
5770
5780
5790
5800
5810
5820
5830
5840
5850
5860
5870
5880
5890
5900
5910
5920
5930
5940
5950
5960
5970
5980
5990
6000
6010
6020
6030
6040
6050
6060
6070
6080
6090
6100
6110
6120
6130
6140
6150
6160
6170
6180
6190
6200
6210
6220
6230
6240
6250
6260
6270
6280
6290
6300
6310
6320
6330
6340
6350
6360
6370
6380
6390
6400
6410
6420
6430
6440
6450
6460
6470
6480
6490
6500
6510
6520
6530
6540
6550
6560
6570
6580
6590
6600
6610
6620
6630
6640
6650
6660
6670
6680
6690
6700
6710
6720
6730
6740
6750
6760
6770
6780
6790
6800
6810
6820
6830
6840
6850
6860
6870
6880
6890
6900
6910
6920
6930
6940
6950
6960
6970
6980
6990
7000
7010
7020
7030
7040
7050
7060
7070
7080
7090
7100
7110
7120
7130
7140
7150
7160
7170
7180
7190
7200
7210
7220
7230
7240
7250
7260
7270
7280
7290
7300
7310
7320
7330
7340
7350
7360
7370
7380
7390
7400
7410
7420
7430
7440
7450
7460
7470
7480
7490
7500
7510
7520
7530
7540
7550
7560
7570
7580
7590
7600
7610
7620
7630
7640
7650
7660
7670
7680
7690
7700
7710
7720
7730
7740
7750
7760
7770
7780
7790
7800
7810
7820
7830
7840
7850
7860
7870
7880
7890
7900
7910
7920
7930
7940
7950
7960
7970
7980
7990
8000
8010
8020
8030
8040
8050
8060
8070
8080
8090
8100
8110
8120
8130
8140
8150
8160
8170
8180
8190
8200
8210
8220
8230
8240
8250
8260
8270
8280
8290
8300
8310
8320
8330
8340
8350
8360
8370
8380
8390
8400
8410
8420
8430
8440
8450
8460
8470
8480
8490
8500
8510
8520
8530
8540
8550
8560
8570
8580
8590
8600
8610
8620
8630
8640
8650
8660
8670
8680
8690
8700
8710
8720
8730
8740
8750
8760
8770
8780
8790
8800
8810
8820
8830
8840
8850
8860
8870
8880
8890
8900
8910
8920
8930
8940
8950
8960
8970
8980
8990
9000
9010
9020
9030
9040
9050
9060
9070
9080
9090
9100
9110
9120
9130
9140
9150
9160
9170
9180
9190
9200
9210
9220
9230
9240
9250
9260
9270
9280
9290
9300
9310
9320
9330
9340
9350
9360
9370
9380
9390
9400
9410
9420
9430
9440
9450
9460
9470
9480
9490
9500
9510
9520
9530
9540
9550
9560
9570
9580
9590
9600
9610
9620
9630
9640
9650
9660
9670
9680
9690
9700
9710
9720
9730
9740
9750
9760
9770
9780
9790
9800
9810
9820
9830
9840
9850
9860
9870
9880
9890
9900
9910
9920
9930
9940
9950
9960
9970
9980
9990
10000
10010
10020
10030
10040
10050
10060
10070
10080
10090
10100
10110
10120
10130
10140
10150
10160
10170
10180
10190
10200
10210
10220
10230
10240
10250
10260
10270
10280
10290
10300
10310
10320
10330
10340
10350
10360
10370
10380
10390
10400
10410
10420
10430
10440
10450
10460
10470
10480
10490
10500
10510
10520
10530
10540
10550
10560
10570
10580
10590
10600
10610
10620
10630
10640
10650
10660
10670
10680
10690
10700
10710
10720
10730
10740
10750
10760
10770
10780
10790
10800
10810
10820
10830
10840
10850
10860
10870
10880
10890
10900
10910
10920
10930
10940
10950
10960
10970
10980
10990
11000
11010
11020
11030
11040
11050
11060
11070
11080
11090
11100
11110
11120
11130
11140
11150
11160
11170
11180
11190
11200
11210
11220
11230
11240
11250
11260
11270
11280
11290
11300
11310
11320
11330
11340
11350
11360
11370
11380
11390
11400
11410
11420
11430
11440
11450
11460
11470
11480
11490
11500
11510
11520
11530
11540
11550
11560
11570
11580
11590
11600
11610
11620
11630
11640
11650
11660
11670
11680
11690
11700
11710
11720
11730
11740
11750
11760
11770
11780
11790
11800
11810
11820
11830
11840
11850
11860
11870
11880
11890
11900
11910
11920
11930
11940
11950
11960
11970
11980
11990
12000
12010
12020
12030
12040
12050
12060
12070
12080
12090
12100
12110
12120
12130
12140
12150
12160
12170
12180
12190
12200
12210
12220
12230
12240
12250
12260
12270
12280
12290
12300
12310
12320
12330
12340
12350
12360
12370
12380
12390
12400
12410
12420
12430
12440
12450
12460
12470
12480
12490
12500
12510
12520
12530
12540
12550
12560
12570
12580
12590
12600
12610
12620
12630
12640
12650
12660
12670
12680
12690
12700
12710
12720
12730
12740
12750
12760
12770
12780
12790
12800
12810
12820
12830
12840
12850
12860
12870
12880
12890
12900
12910
12920
12930
12940
12950
12960
12970
12980
12990
13000
13010
13020
13030
13040
13050
13060
13070
13080
13090
13100
13110
13120
13130
13140
13150
13160
13170
13180
13190
13200
13210
13220
13230
13240
13250
13260
13270
13280
13290
13300
13310
13320
13330
13340
13350
13360
13370
13380
13390
13400
13410
13420
13430
13440
13450
13460
13470
13480
13490
13500
13510
13520
13530
13540
13550
13560
13570
13580
13590
13600
13610
13620
13630
13640
13650
13660
13670
13680
13690
13700
13710
13720
13730
13740
13750
13760
13770
13780
13790
13800
13810
13820
13830
13840
13850
13860
13870
13880
13890
13900
13910
13920
13930
13940
13950
13960
13970
13980
13990
14000
14010
14020
14030
14040
14050
14060
14070
14080
14090
14100
14110
14120
14130
14140
14150
14160
14170
14180
14190
14200
14210
14220
14230
14240
14250
14260
14270
14280
14290
14300
14310
14320
14330
14340
14350
14360
14370
14380
14390
14400
14410
14420
14430
14440
14450
14460
14470
14480
14490
14500
14510
14520
14530
14540
14550
14560
14570
14580
14590
14600
14610
14620
14630
14640
14650
14660
14670
14680
14690
14700
14710
14720
14730
14740
14750
14760
14770
14780
14790
14800
14810
14820
14830
14840
14850
14860
14870
14880
14890
14900
14910
14920
14930
14940
14950
14960
14970
14980
14990
15000
15010
15020
15030
15040
15050
15060
15070
15080
15090
15100
15110
15120
15130
15140
15150
15160
15170
15180
15190
15200
15210
15220
15230
15240
15250
15260
15270
15280
15290
15300
15310
15320
15330
15340
15350
15360
15370
15380
15390
15400
15410
15420
15430
15440
15450
15460
15470
15480
15490
15500
15510
15520
15530
15540
15550
15560
15570
15580
15590
15600
15610
15620
15630
15640
15650
15

14/14

2 amino acid linker

GAA TTC CTG TGC ATG AAA GCA CTG CTA CTC TTC AGC ATT ATT CCA GTC
 glu phe leu cys met lys ala leu leu leu phe ser ile ile pro val 801
 AAC TAC ATC AAG GAA GAT CGC ATC ATT GCA TGC AAA AGA AAA AAT
 asn tyr ile lys glu leu asp arg ile ala cys lys arg lys asn 831
 CTG GAT TCT GTG CAG CCT ATT GCA AGA GAG CTG CAT CAA TTC ACT TTT
 leu asp ser val gln pro ile ala arg glu leu his gln phe thr phe 861
 CAA ATG ATG GCA GAG ATC ATC TCT GTG CAA GTC CCC AAG ATC CTT TCT
 glu met met ala glu ile ile ser val gln val pro lys ile leu ser 891
 CCA ATT TTT GAT GAA CTT CCA ATG AAC TAC ATC AAG GAA GAT CGC ATC ATT GCA TGC AAA AGA AAA AAT
 arg met asn tyr ile lys glu leu asp arg ile ala cys lys arg lys asn 831
 CAC CTC CTA ATG GTC AGC GTC CAC TTT CCT GAA ATG ATG GCA GAG ATC ATC TCT GTG CAA GTC CCC AAG ATC CTT TCT
 asp leu leu ile lys ser his met val ser val ser val gln phe thr phe 861
 GCG AAA GTC AAG CCC ATC ATC TAT TTC CAC ACA CAG TGA
 gly lys val lys pro ile tyr phe his thr gln ...

Total amino acids: 323 + 2 + 117 = 442

FIGURE 8B

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US89/01236

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(4): C07H 21/04, C12N 1/20, C12N 15/00, C07K 13/00

U.S. Cl.: 535/27, 435/252.3, 240.2, 320.5, 7, 530/350, 387

II. FIELDS SEARCHED

Minimum Documentation Searched ⁷

Classification System	Classification Symbols
U.S.	535/27 435/68, 172.3, 240.2, 252.3, 320.5,

Documentation Searched other than Minimum Documentation
to the Extent that such Documents are Included in the Fields Searched ⁸

DATABASES: CHEMICAL ABSTRACTS ONLINE (FILE CA, 1967-1989;
FILE BIOSIS, 1969-1989), USPTO AUTOMATED PATENT SYSTEM
(FILE USPAT, 1975-1988). SEE ATTACHMENT FOR SEARCH TERMS.

III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹

Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
<u>X</u> , P <u>Y</u>	Science (Washington, USA), Volume 240, Issued April 1988, Lubahn et al., "Cloning of human androgen receptor complementary DNA and localization to the X chromosome", pages 327-330, see the entire document.	1-3, 5, <u>10-13</u> 6-8
<u>X</u> , P <u>Y</u>	Science (Washington, USA), Volume 240, Issued April 1988, Chang et al., "Molecular cloning of human and rat complementary DNA encoding androgen receptor", pages 324-326, see the entire document	1-3, 5, 8, <u>10-13</u> 6, 7

^{*} Special categories of cited documents: ¹⁰

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international
filing date

"L" document which may throw doubts on priority claim(s) or
which is cited to establish the publication date of another
citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or
other means

"P" document published prior to the international filing date but
later than the priority date claimed

"T" later document published after the international filing date
or priority date and not in conflict with the application but
cited to understand the principle or theory underlying the
invention

"X" document of particular relevance; the claimed invention
cannot be considered novel or cannot be considered to
involve an inventive step

"Y" document of particular relevance; the claimed invention
cannot be considered to involve an inventive step when the
document is combined with one or more other such docu-
ments, such combination being obvious to a person skilled
in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search
17 June 1989

Date of Mailing of this International Search Report

24 JUL 1989

International Searching Authority
ISA/US

Signature of Authorized Officer *Jasemine C. Chambers*
JASEMINE C. CHAMBERS

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this International application as follows:

I. Claims 1-3, 5-8, 10-13, 30 and 32, drawn to androgen receptor DNA, plasmid, cell and method of use of DNA; Class 435, subclasses 6, 240.2, 252.3 and 320, and Class 536, subclass 27. See attachment.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

1-3, 5-8, 10-13, 30 and 32. Telephone practice.

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

☐ The additional search fees were accompanied by applicant's protest.

☐ No protest accompanied the payment of additional search fees.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	Cold Spring Harbor Symposia on Quantitative Biology, Cold Spring Harbor Laboratory (New York, USA), Volume LI, Published 1986, Mullis et al., "Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction", pages 263-273, see the entire document.	30, 32
Y	Nature (London, UK), Volume 324, Issued November 1986, Saiki et al., "Analysis of enzymatically amplified B-globin and HLA-DQ DNA with allele-specific oligonucleotide probes", pages 163-166, see the entire document.	30, 32
Y, P	US, A, 4,800,159 (MULLIS et al.) 24 JANUARY 1989, see the entire document.	30, 32
X, P	Biochemical and Biophysical Research Communications, Academic Press (Orlando, USA), Volume 153, Issued May 1988, Trapman et al., "Cloning, Structure and expression of a cDNA encoding the human androgen receptor", pages 241-248, see the entire document.	1-3, 5, 8, <u>10-13</u> 6, 7
$\frac{X, P}{Y}$	Proceedings of the National Academy of Sciences, USA (Washington, USA), volume 85, Issued October 1988, Chang et al., "Structural analysis of complementary DNA and amino acid sequences of human and rat androgen receptors", pages 7211-7215, see the entire document.	1-3, 5, 8, <u>10-13</u> 6, 7
$\frac{X}{Y}$	Journal of Endocrinological Investigation (Milan, Italy), Volume 10, Supplement 2, Published 1987, Govindan et al., "Cloning of the human androgen receptor cDNA", page 63, see the entire abstract.	1-3, 5, <u>10-13</u> 6-8
$\frac{X, P}{Y}$	Progress in Cancer Research and Therapy, Raven Press (New York, USA), Volume 35, Issued July 1988, Govindan et al., "Cloning of the human androgen receptor cDNA", pages 49-54, see the entire document.	1-3, 5, <u>10-13</u> 6-8

This Page Blank (uspto)